arXiv:1503.04515v5 [math-ph] 10 Nov 2016

LAX PAIRS OF DISCRETE PAINLEVÉ EQUATIONS: $(A_2+A_1)^{(1)}$ CASE

NALINI JOSHI AND NOBUTAKA NAKAZONO

Abstract. In this paper, we provide a comprehensive method for constructing Lax pairs of discrete Painlevé equations by using a reduced hypercube structure. In particular, we consider the $A_5^{(1)}$-surface q-Painlevé system which has the affine Weyl group symmetry of type $(A_2+A_1)^{(1)}$. Two new Lax pairs are found.

1. Introduction

The purpose of this paper is to provide a comprehensive method for constructing Lax pairs of discrete Painlevé equations by using a reduced hypercube structure arising from ω-lattices, composed from the τ functions of discrete Painlevé equations (see [30, 31] for details). The term "reduced hypercube structure" is used to describe periodic reduction of lattices obtained from multi-dimensionally consistent hypercubes (see §1.3). As an example, we demonstrate the constructions of Lax pairs of the q-Painlevé equations (1.1).

Our previous work [29, 30] brought together a lattice in higher dimensions with τ-function theory, and showed how a geometric reduction provided an ω-lattice leading to $A_5^{(1)}$-surface q-Painlevé equations. Here, we show how this perspective enables us to systematically construct the Lax pairs for any discrete Painlevé equations on the $A_5^{(1)}$-surface. Note that in [54] Sakai classified the discrete Painlevé equations into 22 surface types according to the configuration of the base points (i.e. points where the system is ill defined because it approaches $0/0$) as the following:

Discrete type	Type of surface
Elliptic	$A_0^{(1)}$
Multiplicative	$A_0^{(1)}, \dots, A_8^{(1)}, A_7^{(1)\prime}$
Additive	$A_0^{(1)}, A_1^{(1)}, A_2^{(1)}, D_4^{(1)}, \dots, D_8^{(1)}, E_6^{(1)}, E_7^{(1)}, E_8^{(1)}$

There are possibly infinitely many discrete Painlevé equations on each surface. In this paper, we use the collective term "$A_5^{(1)}$-surface q-Painlevé equations" for discrete Painlevé equations on the multiplicative $A_5^{(1)}$-surface.

Our Lax pairs for the q-Painlevé IV equation (1.1a) and q-Painlevé III equation (1.1b) are new, while the one for the q-Painlevé II equation (1.1c) coincides with that provided in [21]. The spectral part of these Lax pairs is more regular than those from other known Lax pairs. More specifically, it satisfies Carmichael's hypotheses [12] for existence of solutions around singular points at the origin and infinity. These properties allowed Carmichael to prove existence of solutions in a way that is considered to be the q-analogue of Fuchsian theory.

A well-known example which also satisfies Carmichael's conditions is the Lax pair for the q-Painlevé VI equation (q-P($A_3^{(1)}$)), constructed by Jimbo and Sakai [25]. This was seen as a natural analogue of the Fuchsian nature of the Lax pair for the sixth Painlevé equation [17]. However, the Lax pairs of other q-Painlevé equations [39], obtained by degeneration of the Lax pair for q-P($A_3^{(1)}$), no longer satisfy Carmichael's hypotheses: the coefficient matrix of either the highest degree or the lowest degree term in the spectral variable is

---

2010 *Mathematics Subject Classification.* 33E17, 37K10, 39A13, 39A14.

Key words and phrases. Discrete Painlevé equation; ABS equation; Lax pair; τ function; affine Weyl group.

singular. Our results suggest that contrary to expectation, such q-Painlevé equations may still have regular Lax pairs. We provide three examples to support this suggestion.

It is not surprising to have multiple Lax pairs for discrete Painlevé equations as multiple ones are known for each of the continuous Painlevé equations [13, 27, 28].

1.1. **$A_5^{(1)}$-surface q-Painlevé equations.** The q-difference equations we study are

$$q\text{-P}_{\text{IV}}:\ \frac{\overline{f}}{abg}=\frac{1+ch(af+1)}{1+af(bg+1)},\quad \frac{\overline{g}}{bch}=\frac{1+af(bg+1)}{1+bg(ch+1)},\quad \frac{\overline{h}}{caf}=\frac{1+bg(ch+1)}{1+ch(af+1)},\tag{1.1a}$$

$$q\text{-P}_{\text{III}}:\ \overline{g}g=\frac{a(1+tf)}{f(t+f)},\quad \overline{f}f=\frac{a(1+bt\overline{g})}{\overline{g}(bt+\overline{g})},\tag{1.1b}$$

$$q\text{-P}_{\text{II}}:\ \widetilde{f}\underset{\sim}{f}=\frac{a(1+tf)}{f(t+f)},\tag{1.1c}$$

where $t, a, b, c, q, p \in \mathbb{C}^*$ and

$$f=f(t),\quad g=g(t),\quad h=h(t),\quad \overline{f}=f(qt),\quad \overline{g}=g(qt),\quad \overline{h}=h(qt),\tag{1.2a}$$

$$\widetilde{f}=f(pt),\quad \underset{\sim}{f}=f(p^{-1}t).\tag{1.2b}$$

In the case of q-P$_{\text{IV}}$, we have the following conditions:

$$fgh=t^2,\quad abc=q.\tag{1.3}$$

We note that q-P$_{\text{IV}}$, q-P$_{\text{III}}$ and q-P$_{\text{II}}$ are known as a q-discrete analogue of the Painlevé IV equation [36], that of the Painlevé III equation [38, 54] and of the Painlevé II equation [53], respectively.

Remark 1.1. *It is known that q-P$_{\text{III}}$ (1.1b) can be reduced to q-P$_{\text{II}}$ (1.1c) by projective reduction [34]:*

$$b=p,\quad q=p^2,\quad g=\underset{\sim}{f}.\tag{1.4}$$

In this sense, q-P$_{\text{II}}$ is often described as the scalar form of q-P$_{\text{III}}$. Although the projective reduction is a simple specialization of the parameters at the level of the equation, the resulting equations have different type of hypergeometric solutions (see [33, 34] and references therein). In this paper, we also show that they have different Lax pairs but share the same spectral linear problem (see §1.2).

1.2. **Main results.** Our main result shows that Equations (1.1) share one spectral linear problem, which takes a factorized form

$$\phi(qx)=A\phi(x),\tag{1.5}$$

where

$$A=\begin{pmatrix}-\dfrac{\mathrm{i}q\lambda}{f_2}x & 1\\ -1 & -\dfrac{\mathrm{i}qf_2}{\lambda}x\end{pmatrix}\begin{pmatrix}-\dfrac{\mathrm{i}\lambda a_0a_2}{f_0}x & 1\\ -1 & -\dfrac{\mathrm{i}a_0a_2f_0}{\lambda}x\end{pmatrix}\begin{pmatrix}-\dfrac{\mathrm{i}\lambda a_0}{f_1}x & 1\\ -1 & -\dfrac{\mathrm{i}a_0f_1}{\lambda}x\end{pmatrix}.\tag{1.6}$$

Here, the non-zero complex parameters a_i, $i=0,1,2$, λ and q and the variables f_i, $i=0,1,2$, satisfy

$$a_0a_1a_2=q,\quad f_0f_1f_2=\lambda^2.\tag{1.7}$$

However, the deformation problem in the Lax pairs differs for different cases in Equations (1.1). Let $\hat{T}_{\text{IV}}$, $\hat{T}_{\text{III}}$ and $\hat{T}_{\text{SIII}}$ be deformation operators whose actions on the parameters a_i, $i=0,\dots,2$, λ and q are given by

$$\hat{T}_{\text{IV}}:(a_0,a_1,a_2,\lambda,q)\mapsto(a_0,a_1,a_2,q\lambda,q),\tag{1.8a}$$

$$\hat{T}_{\text{III}}:(a_0,a_1,a_2,\lambda,q)\mapsto(qa_0,q^{-1}a_1,a_2,\lambda,q),\tag{1.8b}$$

$$\hat{T}_{\text{SIII}}:(a_0,a_1,a_2,\lambda,q)\mapsto(a_0a_2,q^{-1}a_1a_2,q{a_2}^{-1},\lambda,q),\tag{1.8c}$$

while those on the spectral parameter x and the wave function $\phi=\phi(x)$ are given by

$$\hat{T}_{\mathrm{IV}}(x)=\hat{T}_{\mathrm{III}}(x)=\hat{T}_{\mathrm{SIII}}(x)=x, \tag{1.9a}$$

$$\hat{T}_{\mathrm{IV}}(\phi)=B_{\mathrm{IV}}\,\phi,\quad \hat{T}_{\mathrm{III}}(\phi)=B_{\mathrm{III}}\,\phi,\quad \hat{T}_{\mathrm{SIII}}(\phi)=B_{\mathrm{SIII}}\,\phi, \tag{1.9b}$$

where

$$B_{\mathrm{IV}}=\begin{pmatrix} \dfrac{\mathrm{i}(q\lambda^2-1)f_2}{\lambda(1+a_1(1+a_2f_2)f_1)}x & -1\\ 1 & 0\end{pmatrix}, \tag{1.10a}$$

$$B_{\mathrm{III}}=\begin{pmatrix} -\dfrac{\mathrm{i}\lambda a_0a_2}{f_0}x & 1\\ -1 & -\dfrac{\mathrm{i}a_0a_2f_0}{\lambda}x\end{pmatrix}\begin{pmatrix} -\dfrac{\mathrm{i}\lambda a_0}{f_1}x & 1\\ -1 & -\dfrac{\mathrm{i}a_0f_1}{\lambda}x\end{pmatrix}, \tag{1.10b}$$

$$B_{\mathrm{SIII}}=\begin{pmatrix} -\dfrac{\mathrm{i}\lambda a_0}{f_1}x & 1\\ -1 & -\dfrac{\mathrm{i}a_0f_1}{\lambda}x\end{pmatrix}. \tag{1.10c}$$

The subscripts IV, III, SIII label the deformation operators and matrices corresponding to q-P$_{\mathrm{IV}}$ (1.1a), q-P$_{\mathrm{III}}$ (1.1b), the scalar form of q-P$_{\mathrm{III}}$ (1.1c), respectively. Equations (1.8) and (1.9) provide us with the deformation of the spectral problem.

Theorem 1.2. *The compatibility conditions of the linear equation* (1.5) *with the operators* $\hat{T}_{\mathrm{IV}}$, $\hat{T}_{\mathrm{III}}$ *and* $\hat{T}_{\mathrm{SIII}}$:

$$\hat{T}_{\mathrm{IV}}(A)B_{\mathrm{IV}}=B_{\mathrm{IV}}(qx)A,\quad \hat{T}_{\mathrm{III}}(A)B_{\mathrm{III}}=B_{\mathrm{III}}(qx)A,\quad \hat{T}_{\mathrm{SIII}}(A)B_{\mathrm{SIII}}=B_{\mathrm{SIII}}(qx)A, \tag{1.11}$$

are equivalent to

$$\begin{cases} \dfrac{\hat{T}_{\mathrm{IV}}(f_0)}{a_0a_1f_1}=\dfrac{1+a_2f_2(a_0f_0+1)}{1+a_0f_0(a_1f_1+1)},\\ \dfrac{\hat{T}_{\mathrm{IV}}(f_1)}{a_1a_2f_2}=\dfrac{1+a_0f_0(a_1f_1+1)}{1+a_1f_1(a_2f_2+1)},\\ \dfrac{\hat{T}_{\mathrm{IV}}(f_2)}{a_2a_0f_0}=\dfrac{1+a_1f_1(a_2f_2+1)}{1+a_2f_2(a_0f_0+1)}, \end{cases} \tag{1.12a}$$

$$\hat{T}_{\mathrm{III}}(f_1)f_1=\frac{\lambda^2(1+a_0f_0)}{f_0(a_0+f_0)},\quad \hat{T}_{\mathrm{III}}(f_0)\hat{T}_{\mathrm{III}}(f_1)=\frac{\lambda^2(1+a_0a_2\hat{T}_{\mathrm{III}}(f_1))}{f_0(a_0a_2+\hat{T}_{\mathrm{III}}(f_1))}, \tag{1.12b}$$

$$\hat{T}_{\mathrm{SIII}}(f_1)=f_0,\quad \hat{T}_{\mathrm{SIII}}(f_0)f_1=\frac{\lambda^2(1+a_0f_0)}{f_0(a_0+f_0)}, \tag{1.12c}$$

respectively.

This theorem is proven in §2.4. The actions (1.8) and (1.12) correspond to the q-Painlevé equations (1.1) as explained in the following remark.

Remark 1.3. *Equations* (1.12a) *and* (1.12b) *are equivalent to* q-P$_{\mathrm{IV}}$ (1.1a) *and* q-P$_{\mathrm{III}}$ (1.1b) *by the following correspondences:*

$$\bar{\ }=\hat{T}_{\mathrm{IV}},\quad a=a_0,\quad b=a_1,\quad c=a_2,\quad t=\lambda,\quad f=f_0,\quad g=f_1,\quad h=f_2, \tag{1.13a}$$

$$\bar{\ }=\hat{T}_{\mathrm{III}},\quad a=\lambda^2,\quad b=a_2,\quad t=a_0,\quad f=f_0,\quad g=f_1, \tag{1.13b}$$

respectively. Moreover, letting

$$a_2=q^{1/2}, \tag{1.14}$$

and setting

$$\tilde{\ }=\hat{T}_{\mathrm{SIII}},\quad t=a_0,\quad f=f_0, \tag{1.15}$$

we obtain q-P$_{\mathrm{II}}$ (1.1c) *from Equation* (1.12c).

1.3. **Background.** Discrete Painlevé equations are nonlinear ordinary difference equations of second order, which include discrete analogues of the six Painlevé equations: $\mathrm{P_I}$, …, $\mathrm{P_{VI}}$. The geometric classification of discrete Painlevé equations, based on types of rational surfaces connected to affine Weyl groups, is well known [54]. Together with the Painlevé equations, they are now regarded as one of the most important classes of equations in the theory of integrable systems (see, e.g., [18]).

In [1, 2, 8–10], Adler-Bobenko-Suris (ABS) and Boll classified polynomials P, say, of four variables into eleven types: Q4, Q3, Q2, Q1, H3, H2, H1, D4, D3, D2, D1. The first four types, the next three types and the last four types are collectively called Q-, H^4- and H^6-types, respectively. The resulting polynomial P satisfies the following properties.

(1) Linearity: P is linear in each argument, i.e., it has the following form:

$$P(x_1,x_2,x_3,x_4)=A_1x_1x_2x_3x_4+\cdots+A_{16}, \tag{1.16}$$

where coefficients A_i are complex parameters.

(2) 3D consistency and tetrahedron property: There exist a further seven polynomials of four variables: $P^{(i)}$, $i=1,\dots,7$, which satisfy property **(1)** and a cube C on whose six faces the following equations are assigned

$$P(x_0,x_1,x_2,x_{12})=0, \qquad P^{(1)}(x_0,x_2,x_3,x_{23})=0, \tag{1.17a}$$

$$P^{(2)}(x_0,x_3,x_1,x_{31})=0, \qquad P^{(3)}(x_3,x_{31},x_{23},x_{123})=0, \tag{1.17b}$$

$$P^{(4)}(x_1,x_{12},x_{31},x_{123})=0, \qquad P^{(5)}(x_2,x_{23},x_{12},x_{123})=0, \tag{1.17c}$$

where the eight variables x_0, …, x_{123} lie on the vertices of the cube, in such a way that x_{123} can be uniquely expressed in terms of the four variables x_0, x_1, x_2, x_3 (*3D consistency*) and moreover the following relations hold (*tetrahedron property*):

$$P^{(6)}(x_0,x_{12},x_{23},x_{31})=0, \quad P^{(7)}(x_1,x_2,x_3,x_{123})=0. \tag{1.18}$$

Since these equations relate the vertices of the quadrilateral on a lattice, they are often called quad-equations or lattice equations.

Some polynomials of ABS type are

$$\begin{aligned}
\mathrm{Q1}:\ &Q1(x_1,x_2,x_3,x_4;\alpha_1,\alpha_2;\epsilon)\\
&\quad=\alpha_1(x_1x_2+x_3x_4)-\alpha_2(x_1x_4+x_2x_3)-(\alpha_1-\alpha_2)(x_1x_3+x_2x_4)+\epsilon\alpha_1\alpha_2(\alpha_1-\alpha_2),\\
\mathrm{H3}:\ &H3(x_1,x_2,x_3,x_4;\alpha_1,\alpha_2;\delta;\epsilon)\\
&\quad=\alpha_1(x_1x_2+x_3x_4)-\alpha_2(x_1x_4+x_2x_3)+({\alpha_1}^2-{\alpha_2}^2)\left(\delta+\frac{\epsilon}{\alpha_1\alpha_2}x_2x_4\right),\\
\mathrm{H1}:\ &H1(x_1,x_2,x_3,x_4;\alpha_1,\alpha_2;\epsilon)=(x_1-x_3)(x_2-x_4)+(\alpha_2-\alpha_1)(1-\epsilon x_2x_4),\\
\mathrm{D4}:\ &D4(x_1,x_2,x_3,x_4;\delta_1,\delta_2,\delta_3)=x_1x_3+x_2x_4+\delta_1x_1x_4+\delta_2x_3x_4+\delta_3,
\end{aligned}$$

where $\alpha_1,\alpha_2\in\mathbb{C}^*$ and $\epsilon,\delta,\delta_1,\delta_2,\delta_3\in\{0,1\}$. Many well known integrable PΔEs arise from assigning a polynomial of ABS type to quadrilaterals in the integer lattice $\mathbb{Z}^2$, for example:

discrete Schwarzian KdV equation [41, 43]:

$$Q1(U,\overline{U},\widehat{\overline{U}},\widehat{U};\alpha,\beta;0)=0 \ \Leftrightarrow\ \frac{(U-\overline{U})(\widehat{U}-\widehat{\overline{U}})}{(U-\widehat{U})(\overline{U}-\widehat{\overline{U}})}=\frac{\alpha}{\beta}; \tag{1.19}$$

lattice modified KdV equation [1, 41, 45]:

$$H3(U,\overline{U},-\widehat{\overline{U}},\widehat{U};\alpha,\beta;0;0)=0 \ \Leftrightarrow\ \frac{\widehat{\overline{U}}}{U}=\frac{\alpha\overline{U}-\beta\widehat{U}}{\alpha\widehat{U}-\beta\overline{U}}; \tag{1.20}$$

lattice potential KdV equation [23, 41]:

$$H1(U,\overline{U},\widehat{\overline{U}},\widehat{U};\alpha,\beta;0)=0 \ \Leftrightarrow\ (U-\widehat{\overline{U}})(\overline{U}-\widehat{U})=\alpha-\beta\,; \tag{1.21}$$

discrete version of Volterra-Kac-van Moerbeke equation [41]:

$$\begin{aligned} &D4(1-(\alpha^{-1}\beta-1)U, \widehat{U}, \overline{U}, -1+(\alpha^{-1}\beta-1)\widehat{\overline{U}}; 0,0,0) = 0 \\ \Leftrightarrow\ &\frac{\widehat{U}}{\overline{U}} = \frac{(\beta-\alpha)U-\alpha}{(\beta-\alpha)\widehat{\overline{U}}-\alpha}, \end{aligned} \tag{1.22}$$

where

$$U = U_{l,m}, \quad \alpha = \alpha_l, \quad \beta = \beta_m, \quad \bar{}: l \to l+1, \quad \hat{}: m \to m+1, \quad l, m \in \mathbb{Z}. \tag{1.23}$$

Throughout this paper, we refer to such PΔEs as ABS equations.

We note that in general a hypercube is said to be multi-dimensionally consistent, if all cubes contained in the hypercube are 3D consistent (see property **(2)** above). Reductions of such ABS equations to ordinary difference equations have been found through several approaches [3, 16, 19, 21, 22, 26, 44, 47, 48]. Our geometric-reduction method [29–32] has shown how to obtain discrete Painlevé equations by studying geometric connections between these and ABS equations.

These equations are called integrable because they arise as compatibility conditions for associated linear problems called Lax pairs. The search for and construction of Lax pairs of discrete Painlevé equations has been a very active research area and the investigations have been carried out through many approaches. Noteworthy approaches include extensions of Birkhoff's study of linear q-difference equations [25, 55, 56], periodic-type reductions from ABS equations or the discrete KP/UC hierarchy [20, 21, 37, 47, 49, 50, 52, 57], extensions of Schlesinger transformations [5, 14, 15], search for linearizable curves in initial-value space [35, 61, 62], Padé approximation or interpolation [24, 40, 46] and the theory of orthogonal polynomials [4, 7, 11, 51, 59, 60].

However, the construction of Lax pairs for each case in the literature has been carried out in different ways for different equations on the same surface. In contrast, in the case we study, q-P$_{\rm IV}$ (1.1a), q-P$_{\rm III}$ (1.1b) and q-P$_{\rm II}$ (1.1c) are all obtained on the $A_5^{(1)}$-surface.

1.4. **Plan of the paper.**

The plan of this paper is as follows. In §2, we construct the Lax pairs of the PΔEs on the 4-dimensional integer lattice. Then, we obtain the Lax pairs of the $A_5^{(1)}$-surface q-Painlevé equations from them through geometric reduction. Some concluding remarks are given in §3.

2. THE LAX PAIRS OF THE $A_5^{(1)}$-SURFACE q-PAINLEVÉ EQUATIONS

2.1. **The PΔEs on the lattice $\mathbb{Z}^4$.**

In this section, we consider the PΔEs on the 4-dimensional integer lattice $\mathbb{Z}^4$.

In the same way that the lattice $\mathbb{Z}^2$ can be constructed by tiling the plane with squares, we construct the lattice $\mathbb{Z}^4$ by tiling it with 4-dimensional hypercubes (i.e. 4-cubes). We obtain PΔEs on the lattice $\mathbb{Z}^4$ in a similar manner to the constructions of the ABS equations (see §1.3). Indeed, assigning the function u and quad-equations of ABS type to the vertices and faces of each 4-cube, we obtain the following system of ABS equations:

$$\frac{u(\boldsymbol{l}+\boldsymbol{\epsilon}_1+\boldsymbol{\epsilon}_2)}{u(\boldsymbol{l})}=\frac{u(\boldsymbol{l}+\boldsymbol{\epsilon}_1)-\dfrac{\beta_{l_2}}{\alpha_{l_1}}u(\boldsymbol{l}+\boldsymbol{\epsilon}_2)}{\dfrac{\beta_{l_2}}{\alpha_{l_1}}u(\boldsymbol{l}+\boldsymbol{\epsilon}_1)-u(\boldsymbol{l}+\boldsymbol{\epsilon}_2)},\quad \frac{u(\boldsymbol{l}+\boldsymbol{\epsilon}_1+\boldsymbol{\epsilon}_4)}{u(\boldsymbol{l})}+\frac{u(\boldsymbol{l}+\boldsymbol{\epsilon}_4)}{u(\boldsymbol{l}+\boldsymbol{\epsilon}_1)}=-\alpha_{l_1}K_{l_4},\tag{2.1a}$$

$$\frac{u(\boldsymbol{l}+\boldsymbol{\epsilon}_2+\boldsymbol{\epsilon}_3)}{u(\boldsymbol{l})}=\frac{u(\boldsymbol{l}+\boldsymbol{\epsilon}_2)-\dfrac{\gamma_{l_3}}{\beta_{l_2}}u(\boldsymbol{l}+\boldsymbol{\epsilon}_3)}{\dfrac{\gamma_{l_3}}{\beta_{l_2}}u(\boldsymbol{l}+\boldsymbol{\epsilon}_2)-u(\boldsymbol{l}+\boldsymbol{\epsilon}_3)},\quad \frac{u(\boldsymbol{l}+\boldsymbol{\epsilon}_2+\boldsymbol{\epsilon}_4)}{u(\boldsymbol{l})}+\frac{u(\boldsymbol{l}+\boldsymbol{\epsilon}_4)}{u(\boldsymbol{l}+\boldsymbol{\epsilon}_2)}=-\beta_{l_2}K_{l_4},\tag{2.1b}$$

$$\frac{u(\boldsymbol{l}+\boldsymbol{\epsilon}_3+\boldsymbol{\epsilon}_1)}{u(\boldsymbol{l})}=\frac{u(\boldsymbol{l}+\boldsymbol{\epsilon}_3)-\dfrac{\alpha_{l_1}}{\gamma_{l_3}}u(\boldsymbol{l}+\boldsymbol{\epsilon}_1)}{\dfrac{\alpha_{l_1}}{\gamma_{l_3}}u(\boldsymbol{l}+\boldsymbol{\epsilon}_3)-u(\boldsymbol{l}+\boldsymbol{\epsilon}_1)},\quad \frac{u(\boldsymbol{l}+\boldsymbol{\epsilon}_3+\boldsymbol{\epsilon}_4)}{u(\boldsymbol{l})}+\frac{u(\boldsymbol{l}+\boldsymbol{\epsilon}_4)}{u(\boldsymbol{l}+\boldsymbol{\epsilon}_3)}=-\gamma_{l_3}K_{l_4},\tag{2.1c}$$

where $\boldsymbol{l}=\sum_{i=1}^{4}l_i\boldsymbol{\epsilon}_i\in\mathbb{Z}^4$. Here, $u(\boldsymbol{l})$ is the function on the lattice $\mathbb{Z}^4$ and $\{\alpha_l\}_{l\in\mathbb{Z}}$, $\{\beta_l\}_{l\in\mathbb{Z}}$, $\{\gamma_l\}_{l\in\mathbb{Z}}$ and $\{K_l\}_{l\in\mathbb{Z}}$ are complex parameters. We note that each left hand sides of the equations (2.1) is called $\mathrm{H3}_{\delta=0}$ (or $\mathrm{H3}_{\delta=0}^{\epsilon=0}$) and each right hand sides of them is called D4 in the ABS classification [1, 2, 8].

In the lattice $\mathbb{Z}^4$, there are four orthogonal directions, which naturally give rise to four translation operators. In our case, these result in actions on the variable $u(\boldsymbol{l})$ and the parameters α_l, β_l, γ_l, K_l and lead to the transformations $\hat{T}_i$, $i=1,\dots,4$, by the following actions:

$$\hat{T}_1:(u(\boldsymbol{l}),\alpha_l,\beta_l,\gamma_l,K_l)\mapsto(u(\boldsymbol{l}+\boldsymbol{\epsilon}_1),\alpha_{l+1},\beta_l,\gamma_l,K_l),\tag{2.2a}$$

$$\hat{T}_2:(u(\boldsymbol{l}),\alpha_l,\beta_l,\gamma_l,K_l)\mapsto(u(\boldsymbol{l}+\boldsymbol{\epsilon}_2),\alpha_l,\beta_{l+1},\gamma_l,K_l),\tag{2.2b}$$

$$\hat{T}_3:(u(\boldsymbol{l}),\alpha_l,\beta_l,\gamma_l,K_l)\mapsto(u(\boldsymbol{l}+\boldsymbol{\epsilon}_3),\alpha_l,\beta_l,\gamma_{l+1},K_l),\tag{2.2c}$$

$$\hat{T}_4:(u(\boldsymbol{l}),\alpha_l,\beta_l,\gamma_l,K_l)\mapsto(u(\boldsymbol{l}+\boldsymbol{\epsilon}_4),\alpha_l,\beta_l,\gamma_l,K_{l+1}).\tag{2.2d}$$

In the two-dimensional slice given by $\hat{T}_2$ and $\hat{T}_3$, we define the diagonal region $\mathcal{R}=\nabla^{(1)}\cup\nabla^{(2)}\subset\mathbb{Z}^4$ where

$$\nabla^{(1)}=\left\{\sum_{i=1}^{4}l_i\boldsymbol{\epsilon}_i\;\middle|\;l_i\in\mathbb{Z},\ l_3=l_2-1\right\},\quad \nabla^{(2)}=\left\{\sum_{i=1}^{4}l_i\boldsymbol{\epsilon}_i\;\middle|\;l_i\in\mathbb{Z},\ l_3=l_2\right\}.\tag{2.3}$$

We also define the action of a staircase path $\hat{R}_1$ (see Figure 1) for one of the discrete Painlevé equations considered below:

$$\hat{R}_1(\boldsymbol{l})=\begin{cases}\boldsymbol{l}-\boldsymbol{\epsilon}_2 & \text{if}\quad \boldsymbol{l}\in\nabla^{(1)},\\ \boldsymbol{l}-\boldsymbol{\epsilon}_3 & \text{if}\quad \boldsymbol{l}\in\nabla^{(2)},\end{cases}\tag{2.4}$$

on the variable $u(\boldsymbol{l})$, $\boldsymbol{l}\in\mathcal{R}$, and the parameters α_l, β_l, γ_l, K_l by

$$\hat{R}_1:(u(\boldsymbol{l}),\alpha_l,\beta_l,\gamma_l,K_l)\mapsto\Big(u(\hat{R}_1(\boldsymbol{l})),\alpha_l,\gamma_{l-1},\beta_l,K_l\Big).\tag{2.5}$$

We define the actions of $\hat{T}_i$, $i=1,\dots,4$, and $\hat{R}_1$ on the infinite extension field of the complex field $\mathbb{C}$, generated by $\{u(\boldsymbol{l})\}_{\boldsymbol{l}\in\mathbb{Z}^4\text{ or }\mathcal{R}}$, $\{\alpha_l\}_{l\in\mathbb{Z}}$, $\{\beta_l\}_{l\in\mathbb{Z}}$, $\{\gamma_l\}_{l\in\mathbb{Z}}$ and $\{K_l\}_{l\in\mathbb{Z}}$, as automorphisms. Henceforth, if a new quantity x is added, we extend the field on which $\hat{T}_i$, $i=1,\dots,4$, and $\hat{R}_1$ act as the automorphisms by adding the generator x. Note that throughout this paper every field is of characteristic zero. Moreover, when the field is generated by $\{x_1,\dots,x_k\}$, a mapping $w\in\langle\hat{T}_1,\dots,\hat{T}_4,\hat{R}_1\rangle$ acts on an arbitrary function $F=F(x_1,\dots,x_k)$ by the following:

$$w(F)=F(w(x_1),\dots,w(x_k)).\tag{2.6}$$



Figure 1. Moving of the mapping $\hat{R}_1$ around the origin $\mathbf{0} \in \mathbb{Z}^4$.

For convenience, throughout this paper we use the following notation for the combined transformation of arbitrary mappings w and w':

$$ww' := w \circ w'. \tag{2.7}$$

2.2. **Lax pair of the system of the PΔEs (2.1).**

In this section, we construct the Lax pair of the system of the PΔEs (2.1) following the method given in [6, 42, 58].

The key to the construction of the Lax pair of the system of the PΔEs (2.1) is to introduce fifth direction: $\mathbb{Z}^4 \ni \boldsymbol{l} \mapsto \boldsymbol{l}+\boldsymbol{\epsilon}_5 \in \mathbb{Z}^5$ arising from the extension of the multi-dimensionally consistent 4-cube to the multi-dimensionally consistent 5-cube and to regard it as a virtual direction. As a result, we obtain the following additional relations:

$$\frac{\overline{u}(\boldsymbol{l}+\boldsymbol{\epsilon}_1)}{u(\boldsymbol{l})} = -\frac{\alpha_{l_1}u(\boldsymbol{l}+\boldsymbol{\epsilon}_1)-\mu\overline{u}(\boldsymbol{l})}{\alpha_{l_1}\overline{u}(\boldsymbol{l})-\mu u(\boldsymbol{l}+\boldsymbol{\epsilon}_1)}, \quad \frac{\overline{u}(\boldsymbol{l}+\boldsymbol{\epsilon}_2)}{u(\boldsymbol{l})} = -\frac{\beta_{l_2}u(\boldsymbol{l}+\boldsymbol{\epsilon}_2)-\mu\overline{u}(\boldsymbol{l})}{\beta_{l_2}\overline{u}(\boldsymbol{l})-\mu u(\boldsymbol{l}+\boldsymbol{\epsilon}_2)}, \tag{2.8a}$$

$$\frac{\overline{u}(\boldsymbol{l}+\boldsymbol{\epsilon}_3)}{u(\boldsymbol{l})} = -\frac{\gamma_{l_3}u(\boldsymbol{l}+\boldsymbol{\epsilon}_3)-\mu\overline{u}(\boldsymbol{l})}{\gamma_{l_3}\overline{u}(\boldsymbol{l})-\mu u(\boldsymbol{l}+\boldsymbol{\epsilon}_3)}, \quad \frac{\overline{u}(\boldsymbol{l}+\boldsymbol{\epsilon}_4)}{u(\boldsymbol{l})} + \frac{u(\boldsymbol{l}+\boldsymbol{\epsilon}_4)}{\overline{u}(\boldsymbol{l})} = -\mu K_{l_4}, \tag{2.8b}$$

where $\overline{u}(\boldsymbol{l}) = u(\boldsymbol{l}+\boldsymbol{\epsilon}_5)$, $\boldsymbol{l} = \sum_{i=1}^{4} l_i\boldsymbol{\epsilon}_i \in \mathbb{Z}^4$ and μ is the additional complex parameter. We distinguish the function $\overline{u}(\boldsymbol{l})$ from $u(\boldsymbol{l})$. Then, each of Equations (2.8) can be regarded as the first order discrete system of Riccati type of the quantity $\overline{u}(\boldsymbol{l})$, which is linearizable. Indeed, substituting

$$\overline{u}(\boldsymbol{l}) = \frac{F(\boldsymbol{l})}{G(\boldsymbol{l})}, \tag{2.9}$$

in Equations (2.8) and dividing them into the numerators and the denominators with the vector $\Psi = \Psi(\boldsymbol{l})$ defined by

$$\Psi = \begin{pmatrix} F(\boldsymbol{l}) \\ G(\boldsymbol{l}) \end{pmatrix}, \tag{2.10}$$

we obtain the following linear systems:

$$\hat{T}_1(\Psi) = \delta^{(1)} \begin{pmatrix} \dfrac{\mu}{\alpha_{l_1}} & -\hat{T}_1(u(\boldsymbol{l})) \\ \dfrac{1}{u(\boldsymbol{l})} & -\dfrac{\mu}{\alpha_{l_1}}\dfrac{\hat{T}_1(u(\boldsymbol{l}))}{u(\boldsymbol{l})} \end{pmatrix}\Psi, \quad \hat{T}_2(\Psi) = \delta^{(2)} \begin{pmatrix} \dfrac{\mu}{\beta_{l_2}} & -\hat{T}_2(u(\boldsymbol{l})) \\ \dfrac{1}{u(\boldsymbol{l})} & -\dfrac{\mu}{\beta_{l_2}}\dfrac{\hat{T}_2(u(\boldsymbol{l}))}{u(\boldsymbol{l})} \end{pmatrix}\Psi, \tag{2.11a}$$

$$\hat{T}_3(\Psi) = \delta^{(3)} \begin{pmatrix} \dfrac{\mu}{\gamma_{l_3}} & -\hat{T}_3(u(\boldsymbol{l})) \\ \dfrac{1}{u(\boldsymbol{l})} & -\dfrac{\mu}{\gamma_{l_3}}\dfrac{\hat{T}_3(u(\boldsymbol{l}))}{u(\boldsymbol{l})} \end{pmatrix}\Psi, \quad \hat{T}_4(\Psi) = \delta^{(4)} \begin{pmatrix} -\mu K_{l_4} & -\hat{T}_4(u(\boldsymbol{l})) \\ \dfrac{1}{u(\boldsymbol{l})} & 0 \end{pmatrix}\Psi, \tag{2.11b}$$

where $\boldsymbol{l} = \sum_{i=1}^{4} l_i \boldsymbol{\epsilon}_i$ and $\delta^{(i)} = \delta^{(i)}(\boldsymbol{l})$, $i = 1, \dots, 4$, are arbitrary decoupling factors. The linear systems (2.11) are the Lax pair of the system of the PΔEs (2.1), that is, the compatibility condition of each pair of the linear systems (2.11) gives one of Equations (2.1). For example, the compatibility condition of $\hat{T}_1$ and $\hat{T}_2$:

$$\hat{T}_1\hat{T}_2(\Psi) = \hat{T}_2\hat{T}_1(\Psi), \tag{2.12}$$

gives the left hand side of Equation (2.1a) and

$$\hat{T}_1(\delta^{(2)})\delta^{(1)} = \hat{T}_2(\delta^{(1)})\delta^{(2)}. \tag{2.13}$$

Note that we define the action of a transformation $w \in \langle \hat{T}_1, \dots, T_4, \hat{R}_1 \rangle$ on the vector Ψ by

$$w(\Psi(\boldsymbol{l})) = \Psi(w(\boldsymbol{l})). \tag{2.14}$$

2.3. **Geometric reduction of the PΔEs (2.1).** In this section, we apply the geometric reduction considered in [29] to the PΔEs (2.1). Moreover, we obtain the q-Painlevé equations (1.12) from the reduced PΔEs (2.20).

Let

$$h_{l_1,l_2,l_3,l_4} = \mathrm{i}^{\log(q^{l_1+l_2+l_3+l_4}\hat{\alpha}\hat{\beta}\hat{\gamma}\lambda)/\log q}(q^{l_4}\lambda)^{\log(q^{l_2}\hat{\beta})/\log q}, \tag{2.15}$$

$$a_0 = q\frac{\hat{\alpha}}{\hat{\gamma}}, \quad a_1 = \frac{\hat{\beta}}{\hat{\alpha}}, \quad a_2 = \frac{\hat{\gamma}}{\hat{\beta}}, \tag{2.16}$$

where $a_0a_1a_2 = q$, $\mathrm{i} = \sqrt{-1}$ and $\hat{\alpha}, \hat{\beta}, \hat{\gamma}, \lambda, q \in \mathbb{C}$ are parameters. By letting

$$u(\boldsymbol{l}) = h_{l_1,l_2,l_3,l_4}\,\omega(\boldsymbol{l}), \tag{2.17}$$

where $\boldsymbol{l} = \sum_{i=1}^{4} l_i\boldsymbol{\epsilon}_i$, and imposing the following periodic condition for $\boldsymbol{l} \in \mathbb{Z}^4$:

$$\omega(\boldsymbol{l} + \boldsymbol{\epsilon}_1 + \boldsymbol{\epsilon}_2 + \boldsymbol{\epsilon}_3) = \omega(\boldsymbol{l}), \tag{2.18}$$

with the following condition of the parameters $\{\alpha_l\}_{l\in\mathbb{Z}}$, $\{\beta_l\}_{l\in\mathbb{Z}}$, $\{\gamma_l\}_{l\in\mathbb{Z}}$ and $\{K_l\}_{l\in\mathbb{Z}}$:

$$\alpha_l = q^l\hat{\alpha}, \quad \beta_l = q^l\hat{\beta}, \quad \gamma_l = q^l\hat{\gamma}, \quad K_l = \frac{q^{2l+1}\lambda^2 - 1}{q^l\lambda}, \tag{2.19}$$

the PΔEs (2.1) are reduced to the following PΔEs:

$$\frac{\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_1+\boldsymbol{\epsilon}_2)}{\omega(\boldsymbol{l})} = \frac{\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_1) - q^{-l_1+l_2+l_4}\lambda a_1\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_2)}{q^{l_4}\lambda\,(q^{l_4}\lambda\,\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_2) - q^{-l_1+l_2}a_1\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_1))}, \tag{2.20a}$$

$$\frac{\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_2+\boldsymbol{\epsilon}_3)}{\omega(\boldsymbol{l})} = \frac{q^{l_4}\lambda\,\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_2) - q^{-l_2+l_3}a_2\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_3)}{q^{l_4}\lambda\,(\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_3) - q^{-l_2+l_3+l_4}\lambda a_2\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_2))}, \tag{2.20b}$$

$$\frac{\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_3+\boldsymbol{\epsilon}_1)}{\omega(\boldsymbol{l})} = \frac{\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_3) - q^{l_1-l_3-1}a_0\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_1)}{\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_1) - q^{l_1-l_3-1}a_0\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_3)}, \tag{2.20c}$$

$$\frac{\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_1+\boldsymbol{\epsilon}_4)}{\omega(\boldsymbol{l})} - \frac{\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_4)}{\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_1)} = \frac{q^{2l_4+1}\lambda^2 - 1}{q^{-l_1+l_2+l_4}a_1\lambda}, \tag{2.20d}$$

$$\frac{\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_2+\boldsymbol{\epsilon}_4)}{\omega(\boldsymbol{l})} - \frac{1}{q^{2l_4+1}\lambda^2}\frac{\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_4)}{\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_2)} = \frac{q^{2l_4+1}\lambda^2 - 1}{q^{2l_4+1}\lambda^2}, \tag{2.20e}$$

$$\frac{\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_3+\boldsymbol{\epsilon}_4)}{\omega(\boldsymbol{l})} - \frac{\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_4)}{\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_3)} = \frac{q^{-l_2+l_3}a_2(q^{2l_4+1}\lambda^2 - 1)}{q^{l_4}\lambda}. \tag{2.20f}$$



Figure 2. $(A_2+A_1)^{(1)}$-lattice

Then, the actions of $\hat{T}_i$, $i=1,\dots,4$, and $\hat{R}_1$ on the ω-function and the parameters a_i, $i=0,1,2$, λ and q are given by

$$\hat{T}_1:(\omega(\boldsymbol{l}),a_0,a_1,a_2,\lambda,q)\mapsto(\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_1),qa_0,q^{-1}a_1,a_2,\lambda,q), \tag{2.21a}$$

$$\hat{T}_2:(\omega(\boldsymbol{l}),a_0,a_1,a_2,\lambda,q)\mapsto(\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_2),a_0,qa_1,q^{-1}a_2,\lambda,q), \tag{2.21b}$$

$$\hat{T}_3:(\omega(\boldsymbol{l}),a_0,a_1,a_2,\lambda,q)\mapsto(\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_3),q^{-1}a_0,a_1,qa_2,\lambda,q), \tag{2.21c}$$

$$\hat{T}_4:(\omega(\boldsymbol{l}),a_0,a_1,a_2,\lambda,q)\mapsto(\omega(\boldsymbol{l}+\boldsymbol{\epsilon}_4),a_0,a_1,a_2,q\lambda,q), \tag{2.21d}$$

$$\hat{R}_1:(\omega(\boldsymbol{l}),a_0,a_1,a_2,\lambda,q)\mapsto(\omega(\hat{R}_1(\boldsymbol{l})),a_0a_2,q^{-1}a_1a_2,q{a_2}^{-1},\lambda,q). \tag{2.21e}$$

Remark 2.1. *If we consider the actions of the transformations $\hat{T}_i$, $i=1,\dots,4$, only on the ω-function and the parameters a_i, $i=0,1,2$, λ and q, then the transformation $\hat{T}_1\hat{T}_2\hat{T}_3$ can be regarded as the identity mapping:*

$$\hat{T}_1\hat{T}_2\hat{T}_3=\mathrm{Id}. \tag{2.22}$$

In this situation, we can proceed as if the reduction acts on the lattice $\mathbb{Z}^4$, that is, the lattice $\mathbb{Z}^4$ is reduced to the $(A_2+A_1)^{(1)}$-lattice (see Figure 2):

$$\mathbb{Z}^4\to\mathbb{Z}^4/\mathbb{Z}(\boldsymbol{\epsilon}_1+\boldsymbol{\epsilon}_2+\boldsymbol{\epsilon}_3). \tag{2.23}$$

The reduction from the lattice $\mathbb{Z}^4$ to the $(A_2+A_1)^{(1)}$-lattice with the PΔEs is referred to as the geometric reduction [29].

The relations (2.20) are equivalent to the essential relations of the ω-lattice of type $A_5^{(1)}$ constructed in [30] with the following correspondences:

$$\omega(\boldsymbol{0})=\omega_0,\quad \hat{T}_i=T_i,\ i=1,\dots,4,\quad \hat{R}_1=R_1. \tag{2.24}$$

Therefore, considering the PΔEs (2.20) is equivalent to considering the ω-lattice. In this case, the ω-lattice is said to have a *reduced hypercube structure*. This reduced hypercube structure turns out to be essential in the construction of Lax pairs for discrete Painlevé equations. Since the q-Painlevé equations (1.12) are discrete dynamical systems on the ω-lattice of type $A_5^{(1)}$ as shown in [30], we can obtain the q-Painlevé equations (1.12) from the ω-function with the essential relations (2.20) as follows. By letting

$$f_0=\frac{\omega(\boldsymbol{\epsilon}_1)}{\omega(\boldsymbol{\epsilon}_1+\boldsymbol{\epsilon}_2)},\quad f_1=\lambda\frac{\omega(\boldsymbol{\epsilon}_1+\boldsymbol{\epsilon}_2)}{\omega(\boldsymbol{0})},\quad f_2=\lambda\frac{\omega(\boldsymbol{0})}{\omega(\boldsymbol{\epsilon}_1)}, \tag{2.25}$$

where $f_0f_1f_2=\lambda^2$, we obtain the q-Painlevé equations (1.12) with the following correspondences:

$$\hat{T}_{\mathrm{IV}}=\hat{T}_4,\quad \hat{T}_{\mathrm{III}}={\hat{T}_3}^{-1}{\hat{T}_2}^{-1},\quad \hat{T}_{\mathrm{SIII}}=\hat{R}_1. \tag{2.26}$$

2.4. **Lax pairs of the $A_5^{(1)}$-surface q-Painlevé equations.** In this section, using the linear systems (2.11) and the connection between the PΔEs (2.1) and the $A_5^{(1)}$-surface q-Painlevé equations via the geometric reduction, we construct the Lax pairs of the q-Painlevé equations (1.12).

We first define the vector ϕ by

$$\Psi(\mathbf{0}) = \begin{pmatrix} h_{0,0,0,0}\,\omega(\mathbf{0}) & 0 \\ 0 & 1 \end{pmatrix} \phi. \tag{2.27}$$

Then, from the linear systems (2.11), we obtain the following linear systems:

$$\hat{T}_1(\phi) = \delta^{(1)} \begin{pmatrix} -\dfrac{\mathrm{i} f_2}{\lambda} x & -1 \\ 1 & -\dfrac{\mathrm{i}\lambda}{f_2} x \end{pmatrix} \phi, \tag{2.28a}$$

$$\hat{T}_2(\phi) = \delta^{(2)} \begin{pmatrix} -\dfrac{\mathrm{i} T_1{}^{-1}(f_0)}{a_1 \lambda} x & -1 \\ 1 & -\dfrac{\mathrm{i}\lambda}{a_1 T_1{}^{-1}(f_0)} x \end{pmatrix} \phi, \tag{2.28b}$$

$$\hat{T}_3(\phi) = \delta^{(3)} \begin{pmatrix} -\dfrac{\mathrm{i} a_0 T_3(f_1)}{q \lambda} x & -1 \\ 1 & -\dfrac{\mathrm{i} a_0 \lambda}{q T_3(f_1)} x \end{pmatrix} \phi, \tag{2.28c}$$

$$\hat{T}_4(\phi) = \delta^{(4)} \begin{pmatrix} \dfrac{\mathrm{i}(q\lambda^2 - 1) f_2}{\lambda\,(1 + a_1(1 + a_2 f_2) f_1)} x & -1 \\ 1 & 0 \end{pmatrix} \phi, \tag{2.28d}$$

$$\hat{R}_1(\phi) = \hat{T}_3{}^{-1}(\phi), \tag{2.28e}$$

where

$$x = \frac{\mu}{\hat{\alpha}}. \tag{2.29}$$

Next, let us define the transformations $\hat{T}_{\mathrm{SP}}$, $\hat{T}_{\mathrm{IV}}$, $\hat{T}_{\mathrm{III}}$ and $\hat{T}_{\mathrm{SIII}}$ by

$$\hat{T}_{\mathrm{SP}} = \hat{T}_3{}^{-1}\hat{T}_2{}^{-1}\hat{T}_1{}^{-1}, \quad \hat{T}_{\mathrm{IV}} = \hat{T}_4, \quad \hat{T}_{\mathrm{III}} = \hat{T}_3{}^{-1}\hat{T}_2{}^{-1}, \quad \hat{T}_{\mathrm{SIII}} = \hat{R}_1. \tag{2.30}$$

The actions of $\hat{T}_{\mathrm{SP}}$, $\hat{T}_{\mathrm{IV}}$, $\hat{T}_{\mathrm{III}}$ and $\hat{T}_{\mathrm{SIII}}$ on the spectral parameter x are given by

$$\hat{T}_{\mathrm{SP}}(x) = qx, \quad \hat{T}_{\mathrm{IV}}(x) = \hat{T}_{\mathrm{III}}(x) = \hat{T}_{\mathrm{SIII}}(x) = x, \tag{2.31}$$

while those on the wave function ϕ are given by the following:

$$\hat{T}_{\mathrm{SP}}(\phi) = \frac{\delta^{(\mathrm{SP})}}{(1 - q^2x^2)(1 - a_0{}^2a_2{}^2x^2)(1 - a_0{}^2x^2)} A\,\phi, \quad \hat{T}_{\mathrm{IV}}(\phi) = \delta^{(\mathrm{IV})} B_{\mathrm{IV}}\,\phi, \tag{2.32a}$$

$$\hat{T}_{\mathrm{III}}(\phi) = \frac{\delta^{(\mathrm{III})}}{(1 - a_0{}^2a_2{}^2x^2)(1 - a_0{}^2x^2)} B_{\mathrm{III}}\,\phi, \quad \hat{T}_{\mathrm{SIII}}(\phi) = \frac{\delta^{(\mathrm{SIII})}}{1 - a_0{}^2x^2} B_{\mathrm{SIII}}\,\phi, \tag{2.32b}$$

where

$$\delta^{(\mathrm{SP})} = \frac{1}{\hat{T}_1{}^{-1}\hat{T}_2{}^{-1}\hat{T}_3{}^{-1}(\delta^{(1)})\hat{T}_2{}^{-1}\hat{T}_3{}^{-1}(\delta^{(2)})\hat{T}_3{}^{-1}(\delta^{(3)})}, \quad \delta^{(\mathrm{IV})} = \delta^{(4)}, \tag{2.33a}$$

$$\delta^{(\mathrm{III})} = \frac{1}{\hat{T}_2{}^{-1}\hat{T}_3{}^{-1}(\delta^{(2)})\hat{T}_3{}^{-1}(\delta^{(3)})}, \quad \delta^{(\mathrm{SIII})} = \frac{1}{\hat{T}_3{}^{-1}(\delta^{(3)})}. \tag{2.33b}$$

Here, the 2×2 matrices A, B_{IV}, B_{III} and B_{SIII} are given by Equations (1.6) and (1.10). Therefore, we finally obtain Theorem 1.2 by the following correspondences:

$$\delta^{(1)} = \frac{1}{1-x^2}, \quad \delta^{(2)} = \frac{1}{1-q^{-2}{a_0}^2{a_2}^2x^2}, \quad \delta^{(3)} = \frac{1}{1-q^{-2}{a_0}^2x^2}, \quad \delta^{(4)} = 1, \tag{2.34}$$

which give

$$\delta^{(\mathrm{SP})} = (1-q^2x^2)(1-{a_0}^2{a_2}^2x^2)(1-{a_0}^2x^2), \quad \delta^{(\mathrm{IV})} = 1, \tag{2.35a}$$

$$\delta^{(\mathrm{III})} = (1-{a_0}^2{a_2}^2x^2)(1-{a_0}^2x^2), \quad \delta^{(\mathrm{SIII})} = 1-{a_0}^2x^2. \tag{2.35b}$$

3. CONCLUDING REMARKS

In this paper, we provided a comprehensive method for constructing Lax pairs of discrete Painlevé equations by using a reduced hypercube structure. As an example, we constructed the Lax pairs of the q-Painlevé equations (1.1). As remarked earlier, the discrete Painlevé equations studied in this paper all share one A-matrix in the Lax pairs.

It is possible that we could broaden the action of the affine Weyl group of type $(A_2 + A_1)^{(1)}$, which is the symmetry group for the $A_5^{(1)}$-surface q-Painlevé equations, by adding the actions on the wave function ϕ and the spectral parameter x. This implies that we can construct Lax pairs of "all" $A_5^{(1)}$-surface q-Painlevé equations, which share one spectral linear problem. We will discuss this possibility in a forthcoming paper. Another interesting future direction of research is to extend our method to other surface types of discrete Painlevé equations.

Acknowledgment. The authors would like to express their sincere thanks to Prof. Y. Yamada and Dr. S. Lobb for inspiring and fruitful discussions. This research was supported by an Australian Laureate Fellowship # FL120100094 and grants # DP130100967 and # DP160101728 from the Australian Research Council.

REFERENCES

[1] V. E. Adler, A. I. Bobenko, and Y. B. Suris. Classification of integrable equations on quad-graphs. The consistency approach. *Comm. Math. Phys.*, 233(3):513–543, 2003.

[2] V. E. Adler, A. I. Bobenko, and Y. B. Suris. Discrete nonlinear hyperbolic equations: classification of integrable cases. *Funktsional. Anal. i Prilozhen.*, 43(1):3–21, 2009.

[3] J. Atkinson, P. Howes, N. Joshi, and N. Nakazono. Geometry of an elliptic difference equation related to Q4. *J. Lond. Math. Soc. (2)*, 93(3):763–784, 2016.

[4] P. Biane. Orthogonal polynomials on the unit circle, q-gamma weights, and discrete Painlevé equations. *Mosc. Math. J.*, 14(1):1–27, 170, 2014.

[5] P. Boalch. Quivers and difference Painlevé equations. In *Groups and symmetries*, volume 47 of *CRM Proc. Lecture Notes*, pages 25–51. Amer. Math. Soc., Providence, RI, 2009.

[6] A. I. Bobenko and Y. B. Suris. Integrable systems on quad-graphs. *Int. Math. Res. Not. IMRN*, (11):573–611, 2002.

[7] L. Boelen and W. Van Assche. Discrete Painlevé equations for recurrence coefficients of semiclassical Laguerre polynomials. *Proc. Amer. Math. Soc.*, 138(4):1317–1331, 2010.

[8] R. Boll. Classification of 3D consistent quad-equations. *J. Nonlinear Math. Phys.*, 18(3):337–365, 2011.

[9] R. Boll. Corrigendum: Classification of 3D consistent quad-equations. *J. Nonlinear Math. Phys.*, 19(4):1292001, 3, 2012.

[10] R. Boll. Classification and Lagrangian Structure of 3D Consistent Quad-Equations. *Doctoral Thesis, Technische Universität Berlin*, submitted August 2012.

[11] A. Borodin and D. Boyarchenko. Distribution of the first particle in discrete orthogonal polynomial ensembles. *Comm. Math. Phys.*, 234(2):287–338, 2003.

[12] R. D. Carmichael. The General Theory of Linear q-Difference Equations. *Amer. J. Math.*, 34(2):147–168, 1912.

[13] B. Dubrovin and M. Mazzocco. Monodromy of certain Painlevé-VI transcendents and reflection groups. *Invent. Math.*, 141(1):55–147, 2000.

[14] A. Dzhamay, H. Sakai, and T. Takenawa. Discrete Schlesinger Transformations, their Hamiltonian Formulation, and Difference Painlevé Equations. *arXiv:1302.2972*.

[15] A. Dzhamay and T. Takenawa. Geometric Analysis of Reductions from Schlesinger Transformations to Difference Painlevé Equations. *arXiv:1408.3778*.

[16] C. M. Field, N. Joshi, and F. W. Nijhoff. q-difference equations of KdV type and Chazy-type second-degree difference equations. *J. Phys. A*, 41(33):332005, 13, 2008.

[17] R. Fuchs. Sur quelques équations différentielles linéaires du second ordre. *Comptes Rendus de l'Académie des Sciences Paris*, 141(1):555–558, 1905.

[18] B. Grammaticos and A. Ramani. Discrete Painlevé equations: a review. In *Discrete integrable systems*, volume 644 of *Lecture Notes in Phys.*, pages 245–321. Springer, Berlin, 2004.

[19] B. Grammaticos, A. Ramani, J. Satsuma, R. Willox, and A. S. Carstea. Reductions of integrable lattices. *J. Nonlinear Math. Phys.*, 12(suppl. 1):363–371, 2005.

[20] M. Hay. Hierarchies of nonlinear integrable q-difference equations from series of Lax pairs. *J. Phys. A*, 40(34):10457–10471, 2007.

[21] M. Hay, J. Hietarinta, N. Joshi, and F. Nijhoff. A Lax pair for a lattice modified KdV equation, reductions to q-Painlevé equations and associated Lax pairs. *J. Phys. A*, 40(2):F61–F73, 2007.

[22] M. Hay, P. Howes, N. Nakazono, and Y. Shi. A systematic approach to reductions of type-Q ABS equations. *J. Phys. A*, 48(9):095201, 24, 2015.

[23] R. Hirota. Nonlinear partial difference equations. I. A difference analogue of the Korteweg-de Vries equation. *J. Phys. Soc. Japan*, 43(4):1424–1433, 1977.

[24] Y. Ikawa. Hypergeometric solutions for the q-Painlevé equation of type $E_6^{(1)}$ by the Padé method. *Lett. Math. Phys.*, 103(7):743–763, 2013.

[25] M. Jimbo and H. Sakai. A q-analog of the sixth Painlevé equation. *Lett. Math. Phys.*, 38(2):145–154, 1996.

[26] N. Joshi, B. Grammaticos, T. Tamizhmani, and A. Ramani. From integrable lattices to non-QRT mappings. *Lett. Math. Phys.*, 78(1):27–37, 2006.

[27] N. Joshi, A. V. Kitaev, and P. A. Treharne. On the linearization of the Painlevé III–VI equations and reductions of the three-wave resonant system. *J. Math. Phys.*, 48(10):103512, 42, 2007.

[28] N. Joshi, A. V. Kitaev, and P. A. Treharne. On the linearization of the first and second Painlevé equations. *J. Phys. A*, 42(5):055208, 18, 2009.

[29] N. Joshi, N. Nakazono, and Y. Shi. Geometric reductions of ABS equations on an n-cube to discrete Painlevé systems. *J. Phys. A*, 47(50):505201, 16, 2014.

[30] N. Joshi, N. Nakazono, and Y. Shi. Lattice equations arising from discrete Painlevé systems. I. $(A_2 + A_1)^{(1)}$ and $(A_1 + A_1')^{(1)}$ cases. *J. Math. Phys.*, 56(9):092705, 25, 2015.

[31] N. Joshi, N. Nakazono, and Y. Shi. Lattice equations arising from discrete Painlevé systems. II. $A_4^{(1)}$ case. *J. Phys. A*, (in press) arXiv:1603.09414.

[32] N. Joshi, N. Nakazono, and Y. Shi. Reflection groups and discrete integrable systems. *Journal of Integrable Systems*, (in press) arXiv:1605.01171.

[33] K. Kajiwara and N. Nakazono. Hypergeometric solutions to the symmetric q-Painlevé equations. *Int. Math. Res. Not. IMRN*, (4):1101–1140, 2015.

[34] K. Kajiwara, N. Nakazono, and T. Tsuda. Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$. *Int. Math. Res. Not. IMRN*, (4):930–966, 2011.

[35] K. Kajiwara, M. Noumi, and Y. Yamada. Geometric Aspects of Painlevé Equations. *arXiv:1509.08186*.

[36] K. Kajiwara, M. Noumi, and Y. Yamada. A study on the fourth q-Painlevé equation. *J. Phys. A*, 34(41):8563–8581, 2001.

[37] K. Kajiwara, M. Noumi, and Y. Yamada. q-Painlevé systems arising from q-KP hierarchy. *Lett. Math. Phys.*, 62(3):259–268, 2002.

[38] M. D. Kruskal, K. M. Tamizhmani, B. Grammaticos, and A. Ramani. Asymmetric discrete Painlevé equations. *Regul. Chaotic Dyn.*, 5(3):273–280, 2000.

[39] M. Murata. Lax forms of the q-Painlevé equations. *J. Phys. A*, 42(11):115201, 17, 2009.

[40] H. Nagao. The Padé interpolation method applied to q-Painlevé equations. *Lett. Math. Phys.*, 105(4):503–521, 2015.

[41] F. Nijhoff and H. Capel. The discrete Korteweg-de Vries equation. *Acta Appl. Math.*, 39(1-3):133–158, 1995. KdV '95 (Amsterdam, 1995).

[42] F. W. Nijhoff. Lax pair for the Adler (lattice Krichever-Novikov) system. *Phys. Lett. A*, 297(1-2):49–58, 2002.

[43] F. W. Nijhoff, H. W. Capel, G. L. Wiersma, and G. R. W. Quispel. Bäcklund transformations and three-dimensional lattice equations. *Phys. Lett. A*, 105(6):267–272, 1984.

[44] F. W. Nijhoff and V. G. Papageorgiou. Similarity reductions of integrable lattices and discrete analogues of the Painlevé II equation. *Phys. Lett. A*, 153(6-7):337–344, 1991.

[45] F. W. Nijhoff, G. R. W. Quispel, and H. W. Capel. Direct linearization of nonlinear difference-difference equations. *Phys. Lett. A*, 97(4):125–128, 1983.

[46] M. Noumi, S. Tsujimoto, and Y. Yamada. Padé interpolation for elliptic Painlevé equation. In *Symmetries, integrable systems and representations*, volume 40 of *Springer Proc. Math. Stat.*, pages 463–482. Springer, Heidelberg, 2013.

[47] C. M. Ormerod. Reductions of lattice mKdV to q-P_{VI}. *Phys. Lett. A*, 376(45):2855–2859, 2012.

[48] C. M. Ormerod. Symmetries and special solutions of reductions of the lattice potential KdV equation. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 10:Paper 002, 19, 2014.

[49] C. M. Ormerod, P. H. van der Kamp, J. Hietarinta, and G. R. W. Quispel. Twisted reductions of integrable lattice equations, and their Lax representations. *Nonlinearity*, 27(6):1367–1390, 2014.

[50] C. M. Ormerod, P. H. van der Kamp, and G. R. W. Quispel. Discrete Painlevé equations and their Lax pairs as reductions of integrable lattice equations. *J. Phys. A*, 46(9):095204, 22, 2013.

[51] C. M. Ormerod, N. S. Witte, and P. J. Forrester. Connection preserving deformations and q-semi-classical orthogonal polynomials. *Nonlinearity*, 24(9):2405–2434, 2011.

[52] V. G. Papageorgiou, F. W. Nijhoff, B. Grammaticos, and A. Ramani. Isomonodromic deformation problems for discrete analogues of Painlevé equations. *Phys. Lett. A*, 164(1):57–64, 1992.

[53] A. Ramani and B. Grammaticos. Discrete Painlevé equations: coalescences, limits and degeneracies. *Phys. A*, 228(1-4):160–171, 1996.

[54] H. Sakai. Rational surfaces associated with affine root systems and geometry of the Painlevé equations. *Comm. Math. Phys.*, 220(1):165–229, 2001.

[55] H. Sakai. A q-analog of the Garnier system. *Funkcial. Ekvac.*, 48(2):273–297, 2005.

[56] H. Sakai. Lax form of the q-Painlevé equation associated with the $A_2^{(1)}$ surface. *J. Phys. A*, 39(39):12203–12210, 2006.

[57] T. Tsuda. On an integrable system of q-difference equations satisfied by the universal characters: its Lax formalism and an application to q-Painlevé equations. *Comm. Math. Phys.*, 293(2):347–359, 2010.

[58] A. Walker. Similarity reductions and integrable lattice equations. *Ph.D. Thesis, University of Leeds*, submitted September 2001.

[59] N. S. Witte. Semiclassical orthogonal polynomial systems on nonuniform lattices, deformations of the Askey table, and analogues of isomonodromy. *Nagoya Math. J.*, 219:127–234, 2015.

[60] N. S. Witte and C. M. Ormerod. Construction of a Lax pair for the $E_6^{(1)}$ q-Painlevé system. *SIGMA Symmetry Integrability Geom. Methods Appl.*, 8:Paper 097, 27, 2012.

[61] Y. Yamada. A Lax formalism for the elliptic difference Painlevé equation. *Symmetry Integrability Geom. Methods Appl.*, 5:Paper 042, 15, 2009.

[62] Y. Yamada. Lax formalism for q-Painlevé equations with affine Weyl group symmetry of type $E_n^{(1)}$. *Int. Math. Res. Not. IMRN*, (17):3823–3838, 2011.

School of Mathematics and Statistics, The University of Sydney, New South Wales 2006, Australia.
E-mail address: `nobua.n1222@gmail.com`